

Humantech's Position

# Physical Risk Factors for Musculoskeletal Disorders



## Introduction

Musculoskeletal disorders (MSDs) are defined as injuries or dysfunctions affecting muscles, bones, nerves, tendons, ligaments, joints, cartilages, and spinal discs. They include sprains, strains, carpal tunnel syndrome (CTS), hernias, and connective tissue injuries of these structures. MSDs are among the leading causes of lost workdays and are associated with a large economic burden according to the American Academy of Orthopedic Surgeons (AAOS).

MSDs account for approximately one-third of all workplace injuries and illnesses in the United States (BLS, 2015). According to the Liberty Mutual Workplace Safety Index (2016), overexertion injuries (a particular type of MSD) are the leading cause of all non-fatal workplace injuries in the United States. Overexertion activities are defined as lifting, lowering, pushing, pulling, carrying, holding, or throwing. In 2013, overexertion injuries accounted for 24.4% of all non-fatal workplace injuries and amounted to \$15.08 billion in direct costs (medical costs and indemnity payments). The indirect costs associated with workplace MSDs (production time lost by the injured employee, fellow workers, and supervisors; spoiled product; unhappy customers; cleanup time; schedule delays; training new employees; overhead costs; legal fees; and an increase in insurance costs) are also significant and can amount to a further 1.1 to 4.5 times that of the direct costs.

This statement provides background for our position on the primary physical risk factors in the workplace that lead to the development of MSDs.

Our position is that forceful and prolonged exertions, awkward postures, and repetition are the primary physical risk factors in the workplace that are associated with the development of MSDs. Two of the risk factors, forceful exertions and awkward postures, seem to play a greater role, and there are complex interactions (interdependence) between certain risk factors that influence the development of MSDs.

## Physical Risk Factors and MSDs

For over a century, an association between physical risk factors in the workplace and MSDs has been observed. In the late 1990s, The National Institute of Occupational Safety and Health (NIOSH) published the most comprehensive compilation of the epidemiological studies on the correlation of exposure to physical factors in the workplace and MSDs (NIOSH, 1997). NIOSH concluded that there is a large body of credible epidemiological research that shows a consistent relationship between certain physical risk factors in the workplace and MSDs. These physical risk factors include repetitive motion, excessive force, and awkward postures (including sustained postures, prolonged sitting, and standing).

Since the 1990s, there have been several excellent epidemiological studies (both case control and cohort studies) that have added to our understanding of the physical risk factors in the workplace that lead to the development of MSDs. These studies have shown a reasonable causal relationship between physical risk factors in the workplace and the development of MSDs for a wide range of body areas and injuries/illnesses (CTS, lateral epicondylitis, tendinitis, low back pain, sciatica, and hip/knee osteoarthritis). Recent research examples of the causal or dose-response relationship between physical risk factors in the workplace and MSDs include the following:

- **Hand/wrist disorders.** Recently, in a large multi-site, cross-sectional cohort study, Fan et. al (2015) found a strong dose-response relationship between jobs requiring high hand force and CTS; however, for jobs requiring either certain wrist postures or repetition, but not both, this same relationship did not exist. Also, Kapellusch et. al (2014) demonstrated that the ACGIH TLV for Hand Activity Level (HAL) is a strong predictor of CTS.



- **Shoulder disorders.** Hoozemans et. al (2014) summarized high-quality cohort studies and showed there is strong evidence that pushing and pulling are related to upper extremity symptoms, specifically for the shoulder.
- **Low back disorders.** Garg et al. (2014) showed a statistically significant exposure-response relationship between both the NIOSH Lifting Equation peak lifting force (PLI) and peak composite lifting index (PCLI) and low back pain.

Overall, these excellent epidemiological studies have added to our understanding, and the primary risk factors in the workplace for each body area are summarized below:

Body area	Biomechanical Risk Factors			
	Awkward postures	High forces	Long durations	High frequency
Neck	•	•	•	•
Low back	•	•	•	•
Upper limb	•	•	•	•
Shoulder		•		•
Elbow/forearm	•	•	•	•
Wrist/hand	•	•	•	•
Lower limb	•	•	•	•

## Interaction of Physical Risk Factors is Key

For many years, it was assumed that each physical risk factor in the workplace functioned as an independent factor, without any interdependence with other risk factors. Recently, Gallagher and Heberger (2013) addressed this assumption. Based on several high-quality studies, they showed that there is a strong interdependence between high force and high repetition across a wide range of joint disorders and symptoms, including low back disorders, carpal tunnel syndrome, hand-wrist tendinitis, wrist discomfort, lateral epicondylitis, shoulder tendinitis, shoulder discomfort, and knee discomfort. To summarize, they found that:

- Tasks that expose individuals to low force, at either a low repetition or high repetition, only nominally impact the risk of joint injury.
- Tasks that expose individuals to high force and low repetition have a low risk of joint injury.
- Tasks that expose individuals to high force and high repetition significantly increase the risk of joint injury.

In addition to finding that risk factors are interdependent, they also showed that certain risk factors, such as forceful exertion, play a greater role in the development of MSDs.

Since awkward postures can alter the force capabilities of a joint, the relationship between force and joint posture also shows that the awkward postures play a more significant part in the development of MSDs. Thus, forceful exertions and awkward postures are two primary, and complimentary, influences on the risk of joint injury.

## Conclusion

Based on strong epidemiological evidence, there is a good understanding of the physical risk factors in the workplace that lead to the development of MSDs for several body areas and joints. Some risk factors, such as forceful exertion and awkward postures, seem to contribute more significantly to the development of MSDs. However, the development of MSDs is influenced by complex relationships among risk factors, meaning there is some interdependence.

An MSDs risk assessment tool should cover all of the main body areas, focus on measuring the primary physical risk factors in the workplace (such as forceful exertions and awkward postures), and should also capture the complex interaction between factors (such as repetition and duration).

## Endorsement

This position statement was accepted by Senior Leadership on October 24, 2016.

## References

- American Academy of Orthopedic Surgeons. 2008. *The Burden of Musculoskeletal Disorders in the United States: Prevalence, Societal, and Economic Cost*. Executive Summary. Rosemount, IL.
- Bureau of Labor Statistics. 2015. *Non-fatal Occupational Injuries and Illnesses Requiring Days Away from Work, 2014*. November 19, 2015.
- da Costa BR and Vieira ER. 2010. *Risk Factors for Work-Related Musculoskeletal Disorders: A Systematic Review of Recent Longitudinal Studies*. American Journal of Industrial Medicine 53: 285-323.
- Fan ZJ, Harris-Adamson C, Geer F, Eisen EA, Hegmann KT, Bao S, Silverstein B, Evanoff B, Dale AM, Thiese MS, Garg A, Kapellusch J, Burt S, Merlino L, and Rempel D. 2015. *Associations between Workplace Factors and Carpal Tunnel Syndrome: A Multi-Site Cross Sectional Study*. American Journal of Industrial Medicine. 58: 509-518.



Garg A, Boda S, Hegmann KT, Moore JS, Kapellusch JM, Bhojar P, Thiese MS, Merryweather A, Decker-Schaefer G, Blosswick D, and Malloy EJ. 2014. *The NIOSH Lifting Equation and Low-Back Pain, Part 1: Association with Low-Back Pain in the BackWorks Prospective Cohort Study*. Human Factors. 56(1) 6-28.

Garg A, Waters T, Kapellusch J, and Karwowski W. 2014b. *Psychophysical Basis for Maximum Pushing and Pulling Forces: A Review and Recommendations*. International Journal of Industrial Ergonomics. 44: 281-291.

Hoozemans MJ, Knelange EB, Frings-Dresen MH, Veeger HE, and Kuijer PP. 2014. *Are pushing and pulling work-related risk factors for upper extremity symptoms? A systematic review of observational studies*. Occupational and Environmental Medicine. 71(11):788-95

Horton, R. (2012). *Understanding Disease, Injury, and Risk*. Lancet. (380):2053-2054.

Kapellusch JM, Gerr FE, Malloy EJ, Garg A, Harris-Adamson C, Bao SS, Burt SE, Dale AM, Eisen AE, Evanoff BA, Hegmann KT, Silverstein BA, Thiese MS and Rempel DM. 2014. *Exposure-Response Relationship for the ACGIH Threshold Limit Value for Hand-Activity Level: Results from a Pooled Data Study of Carpal Tunnel Syndrome*. Scandinavian Journal of Work, Environment, and Health. 40(6): 610-620

Liberty Mutual Research Institute for Safety. 2016. *Liberty Mutual Workplace Safety Index*. Executive Summary. Hopkinton, MA.

National Institute for Occupational Safety and Health (NIOSH). 1997. *Musculoskeletal Disorders and Workplace Factors: A Critical Review of Epidemiologic Evidence for Work-Related Musculoskeletal Disorders of the Neck, Upper Extremity, and Low Back*. DHHS (NIOSH) Publication No. 97-141.

